

# Assessing land cover changes in the French Pyrenees since the 1940s A semi-automatic GEOBIA approach using aerial photographs

D Sheeren, S Ladet, O Ribière, B Raynaud, Martin Paegelow, Thomas Houet

► **To cite this version:**

D Sheeren, S Ladet, O Ribière, B Raynaud, Martin Paegelow, et al.. Assessing land cover changes in the French Pyrenees since the 1940s A semi-automatic GEOBIA approach using aerial photographs. AGILE 2012 : International conference on geographic information science, Apr 2012, Avignon, France. hal-01197132

**HAL Id: hal-01197132**

**<https://hal-univ-tlse2.archives-ouvertes.fr/hal-01197132>**

Submitted on 11 Sep 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Assessing land cover changes in the French Pyrenees since the 1940s


## A semi-automatic GEOBIA approach using aerial photographs



**D. Sheeren, S. Ladet**  
 DYNAFOR Lab.  
 UMR 1201 INP-ENSAT / INRA  
 University of Toulouse  
 BP 32607, 31326 Castanet Tolosan Cedex, France

**O. Ribière, B. Raynaud, M. Paegelow, T. Houet**  
 GEODE Lab.  
 UMR 5602 CNRS / UTM  
 University of Toulouse  
 5 al. Machado, 31058 Toulouse, France

### Introduction




As in many mountain areas in Europe, **agro-pastoral landscapes of the Pyrenees are subject to fast spontaneous reforestation**. Encroachment is mainly related to the abandonment of farmland in combination with rural depopulation and agriculture modernization.

Observed changes of land abandonment or land-use extensification can be similar at the regional scale but the **dynamics at the local scale do not occur homogeneously**, as it is determined by specific conditions (environmental, climatic, socio-economic, anthropogenic).

The **objective** of this work (part of the MODE-RESPYR project) is to **assess the spatial patterns of land cover changes** during the last 70 years in three study sites of the Pyrenees, and to **compare the local dynamics** in order to **observe and to explain similarities and disparities**.

### Study sites




- 1. Davantaygue (76 km<sup>2</sup>)**
  - In the Peripheral area of the Pyrenees National Park
  - 6 municipalities
- 2. Haut-Vicdessos (244 km<sup>2</sup>)**
  - In the Regional Natural Park of Ariège
  - 7 municipalities
- 3. Garrotxes (85 km<sup>2</sup>)**
  - In the Regional Natural Park of the Catalan Pyrenees
  - 5 municipalities

### Material and Method

#### Data


Historical panchromatic aerial photographs with recent true color digital images



- 1. Davantaygue**
  - Number of photographs: 67
  - 6 dates: from 1948 to 2001
  - Scale: 1:15000-1:30000
  - Spatial resolution: 0.5m
  - TIFF (8 or 24 bits)
- 2. Haut-Vicdessos**
  - Number of photographs: 286
  - 7 dates: from 1942 to 2003
  - Scale: 1:25000-1:30000
  - Spatial resolution: 0.5m
  - TIFF (8 or 24 bits)
- 3. Garrotxes**
  - Number of photographs: 104
  - 5 dates: from 1942 to 2000
  - Scale: 1:25000-1:30000
  - Spatial resolution: 0.5m
  - TIFF (8 or 24 bits)


#### Methodological approach

Dependently-generated land cover maps from a GEOgraphic Object-Based Image Analysis




### Results

#### 1. Davantaygue



#### 2. Haut-Vicdessos



### Conclusions

- A similar spatial pattern of land cover changes for the 3 study sites (encroachment at low and medium altitudes) but with **different rates and extents**, and occurring at different periods.
- A spatial pattern highly related to pattern of **farmland abandonment** and depopulation.
- A **generic method** usable for various mountain areas
- Perspectives: developing an automatic classification procedure using very high resolution remotely sensed images for monitoring annual land use and cover changes.

#### 3. Garrotxes

